

# Research Challenge Beef Farm Walk

***“On-farm research to underpin improvements in the carbon footprint of beef production”***

at the farm of:

**Arti Birt**

30 Deerpark Road, Portaferry



**Tuesday 24<sup>th</sup> July, 2012**

## Research to underpin improved production efficiency

Today's farm walk is aimed at providing you with tools and information to help you make improvements with your beef enterprise

### Topics for discussion include:

1. How to the reduce the age at first calving and the impact
2. Pain free performance monitoring
3. Grassland management in good and bad weather
4. The economics of suckler beef production

## Arti Birt, Portaferry

- Farm Area: 290 acres
- 160 suckler cows of Limousin X Charolais X Simmental (3 way cross)
- Majority of progeny finished

### Aims:

- Maximising production efficiency
  - monitoring performance
  - efficient use of labour
  - calving at 24 months of age - ideal body condition
  - 25 acres reseeded annually
  - bulls based on EBV figures



### Recent Farm Developments & Technology include:

- Regular weighing of cattle to monitor growth
- Modifying diets in line with animal performance

## Target weights for rearing replacements

Mature cow weight 650 kg

Bulling weight  
60% mature  
weight at  
14 months


Calving weight  
90% mature  
weight at  
24 months

Age (months)	Weight (kg)	Growth rate (kg/d)
3	110	0.90
6	215	
9	280	
12	330	0.74
14	390	
18	480	
21	532	0.57
24	585	


Key is to monitor performance – online tool being developed to help with this

## Arti Birt, Portaferry


SUMMARY	Spring 2010
Mature cow weight	620 kg
Target weight at 1 <sup>st</sup> calving	555 kg
Target weight at breeding	370 kg
No. of animals	43
Age	15 months
Live weight	447 kg
DLWG achieved	0.89 kg/day
No. of animals	34
Age	20 months
Live weight	540 kg
DLWG achieved	0.82 kg/day



SUMMARY	Spring 2011
Mature cow weight	620 kg
Target weight at 1 <sup>st</sup> calving	555 kg
Target weight at breeding	370 kg
No. of animals	55
Age	11 months
Live weight	342 kg
DLWG achieved	0.89 kg/day
No. of animals	40
Age	13 months
Live weight	389 kg
DLWG achieved	0.90 kg/day



- ◆ Online tool to aid growth monitoring
- ◆ Animal list and ages supplied by APHIS
- ◆ Weights automatically plotted against target



Animal Type: Suckler Herd Replacements

Age at first calving: 24 months

Mature Cow Weight:  kg

Calving Weight:  kg

Animal Tag No	Sex	Breed	Date of Birth	Age (months)	Weight (kg)
UK 9 390002 8274 4	F	Aberdeen-Angus	10/02/2011	17.2	<input type="text" value="400"/>
UK 9 390002 8282 5	F	Charolais	15/02/2011	17.0	<input type="text" value="440"/>
UK 9 390002 8284 7	F	Charolais	20/02/2011	16.9	<input type="text" value="405"/>
UK 9 390002 8286 2	F	Aberdeen-Angus	28/02/2011	16.6	<input type="text" value="395"/>
UK 9 390002 8290 6	F	Aberdeen-Angus	09/03/2011	16.3	<input type="text" value="350"/>
UK 9 390002 8291 7	F	Stabiliser	11/03/2011	16.2	<input type="text" value="300"/>
UK 9 390002 8292 1	F	Charolais	12/03/2011	16.2	<input type="text" value="410"/>
UK 9 390002 8294 3	F	Aberdeen-Angus	14/03/2011	16.1	<input type="text" value="390"/>
UK 9 390002 8295 4	F	Aberdeen-Angus	19/03/2011	16.0	<input type="text" value="305"/>
UK 9 390002 8296 5	F	Charolais	20/03/2011	15.9	<input type="text" value="350"/>
UK 9 390002 8297 6	F	Charolais	22/03/2011	15.9	<input type="text" value="350"/>
UK 9 390002 8300 2	F	Stabiliser	23/03/2011	15.8	<input type="text" value="430"/>
UK 9 390002 8707 3	F	Charolais	10/04/2011	15.2	<input type="text" value="395"/>
UK 9 390002 8708 4	F	Charolais	12/04/2011	15.2	<input type="text" value="410"/>
UK 9 390002 8711 7	F	Stabiliser	22/04/2011	14.9	<input type="text" value="400"/>
UK 9 390002 8710 6	F	Stabiliser	26/04/2011	14.7	<input type="text" value="300"/>

## Bovine Information System (BovIS)

	RCF farms (11/12) <sup>1</sup>	NI average <sup>2</sup>
Age at first calving (months)	24	31
Calving interval (d)	368	400
Calves per cow per year	0.95	0.83
Females not calved (%)	4.0	10.7
% of heifers calved 22-26 months of age	72	18
% of herd calving within 90 days	75	68

<sup>1</sup>Based on four out of the six RCF farms as two had yet to complete the calving season when reports created

<sup>2</sup>Based on approximately 250 Northern Ireland suckler herds

### Tools now available:

- ◆ To benchmark physical and financial performance (CAFRE benchmarking and BovIS)
- ◆ Help producers easily monitor performance (BovIS growth monitoring tool)

Farm	No. calved	Calves born dead/dead within 24 hrs	Calf birth weight (kg)	No. of veterinary assisted calvings
A	33	1	36	2
B	11	0	32	0
C	31	3	37	0
D	16	1 (twin)	42	1
E	10	1	36	0
F	36	4	38	3
Hillsborough	21	1	36	0

**On average 4% of heifers required veterinary assistance at calving**

## CAFRE EXPERIENCE

	2 year calving <sup>1</sup>	Mature cows
Weight at weaning (kg) <sup>2</sup>	576	666
Calf gain (kg/d)	1.01	1.10
200d weight (kg)	245	264
Weaning efficiency <sup>3</sup>	42.5	40.3
Percentage back in calf (%)	93	94

<sup>1</sup> Easy calving sire used

<sup>2</sup> 2yo and 3yo heifers attain similar mature weights

<sup>3</sup> Calf weight at 200 days per 100kg cow weight

- ◆ CAFRE has been successfully calving heifers at 2 years of age since 2007  
2year old heifers are consistently the most efficient age group in the herd

## Performance of RCF producers

RCF project Farm	Age at calving (2011/12)
A	23
B	27 (purchased heifers)
C	25
D	26 (24)
E	27 (25)
F	23
Hillsborough	25

RCF herd age at first calving has decreased by 3.3 months since starting the project

## Improved efficiency of production is key!

### What are the greenhouse gases associated with agriculture?


**Carbon dioxide** – fertiliser, lime, herbicides, pesticides, fuel, electricity, animal feed etc

**Nitrous oxide** – fertiliser application, manure management, crop residues etc

**Methane** – enteric fermentation in rumen, manure management

### What are we doing about it?

- ◆ DEFRA and the devolved administration have funded a large project involving AFBI and research organisations across the UK to improve the accuracy of the national GHG inventory
- ◆ DARD and AgriSearch have commissioned AFBI to investigate GHG reduction strategies and develop a tool to enable producers to calculate emissions from their own farm
- ◆ Greenhouse Gas Reduction Strategy and Action Plan



## Improved efficiency of production is key!

### Reducing the age at first calving


- ◆ Less animals on the farm
- ◆ Less inputs required such as feed and land
- ◆ Increased carcass output per ha
- ◆ Improved cow longevity and fertility

### Other methods to reduce GHG emissions:

- ◆ Reducing the age at slaughter
- ◆ Balanced diet formulation
- ◆ The use of improved beef genetics
- ◆ Minimising animal mortality and morbidity
- ◆ Improved fertility
- ◆ Efficient use of fertiliser nitrogen, clover, legumes

Effect of age at first calving on the number of replacement heifers on the farm (100 cow herd)

Age group	Age at calving (months)	
	24	36
0-12	20	20
12-24	20	20
24-36	0	20



**Improved production efficiency is key!**

## Making the most of grass/grass silage

### 0 – 12 months

- ◆ Weaned at 8 months of age 300 kg (approx)
- ◆ Good grass silage plus 1-2 kg meal/d
- ◆ Early turnout to pasture

Analysis	RCF farmers
Dry matter (%)	27.8
ME (MJ/kg DM)	10.6
D Value	66
Protein (%)	11.7

### 12 – 20 months

- ◆ Good grassland management – rotational grazing
- ◆ Bullied at 60-65% mature weight and CS 3
- ◆ Careful monitoring of weight/CS

Requirements until point of calving	RCF farmers
Total silage fed	5.5 t
Total meal fed	373 kg

### 20 – 24 months

- ◆ Housed on good silage alone with min/vit
- ◆ Careful monitoring of weight/CS

## Importance of high quality grass silage

Silage quality	D –value (% DM)	330 kg continental heifer		515 kg in-calf continental heifer	
		Growth rate from silage alone (kg/d)	Conc required to achieve 0.74 kg/day	Growth rate from silage alone (kg/d)	Conc required to achieve 0.5 kg/day
High	77	0.95	0	1.0	0*
Low	60	0.04	4.5	0.01	3.5
<b>Average</b>	<b>67</b>	<b>0.44</b>	<b>2.0</b>	<b>0.45</b>	<b>0.50</b>

\* Need to restrict intake

◆ High quality grass silage will reduce concentrate requirement

## Industry Analysis – BovIS and LMC data

	Number slaughtered in 2011	Percentage from birth to finish farms	Age at slaughter (months)	Carcass weight (kg)	Conformation grade	Fat class
Bulls	19306	42	17	372	R+	3-
Steers	70744	18	26	369	R=	3=
Heifers	66402	25	25	317	R=	3+

- ◆ Important source of high quality beef – 39% in spec compared to 15% of dairy origin cattle
- ◆ Suckler beef production important role in managing and shaping the countryside

Animal type	No. killed	Carcass weight (kg)	Age at slaughter (months)	Grade	Fat class	% In Spec
Bulls	19	348	15	R+	3-	64.7
Steers	59	368	23	R+	3+	62.3
Heifers	54	309	23	R=	3=	71.4

Animal type	Target daily carcass gain (kg/d)	Daily carcass gain achieved (kg/d)
Bulls	0.73	0.74
Steers	0.49	0.53
Heifers	0.47	0.45

- Key objective on the Birt farm is to produce prime quality carcasses in the most efficient manner
  - ✓ Growth targets and monitoring performance